

Cationic Bis(oxazoline)Cu(II) Lewis Acid Catalysts. Enantioselective Furan Diels-Alder Reaction in the Synthesis of *ent*-Shikimic Acid


David A. Evans* and David M. Barnes

Department of Chemistry & Chemical Biology, Harvard University, Cambridge, MA 02138, USA

Abstract: The highly enantioselective Diels-Alder reaction between acryloyl oxazolidinone and furan, catalyzed by cationic bis(4-*tert*-butyloxazoline)Cu(II) complex **1**, is presented. Though the reaction equilibrates rapidly at -20 °C, reaction at -78 °C permits isolation of the kinetic product mixture. The synthetic utility of the reaction is demonstrated by the conversion of the cycloadduct to *ent*-shikimic acid.
 Copyright © 1996 Published by Elsevier Science Ltd

Recent publications from this laboratory have documented the development of cationic copper(II) complexes as chiral Lewis acids in the Diels-Alder¹ and Mukaiyama aldol² reactions. In particular, complex **1** has been found to be the optimal catalyst for a number of Diels-Alder reactions. In this Letter, we demonstrate the utility of this catalyst system in the reaction of acrylimide **2** with furan to yield cycloadduct **3**, as well as the conversion of the latter to shikimic acid.

Scheme I




Although 7-oxabicyclo[2.2.1]hept-2-enes are useful synthetic intermediates,³ only two examples of catalytic asymmetric Diels-Alder reactions with furan or substituted furans have been reported, one employing α -haloacroleins as the dienophiles,⁴ and the other requiring the more reactive 3-(methylthio)furan as the diene component.⁵ Our investigation of this system began with the investigation of acrylimide **2** with furan in the presence of 5 mol% of complex **1** at -20 °C for 24 h. Cycloadduct **3** was isolated as a 9:1 mixture of *exo* to *endo* diastereomers. Surprisingly, both cycloadducts were found to be racemic. However, when the reaction was terminated after 2.5 h (88% conversion), the *endo*:*exo* ratio had reversed to 66:34, with the major *endo* adduct obtained in 59% ee. These results suggested that at -20 °C, the reaction was equilibrating rapidly to generate a thermodynamic mixture of products. Accordingly, when the reaction was performed at -78 °C, the *endo*:*exo* diastereomer ratio climbed to 80:20, and the *endo* isomer was obtained in 97% ee. Recrystallization of a 20 mmol scale reaction delivered enantiomerically pure **3**, mp 89 °C, $[\alpha]_D^{25} +99.1^\circ$ ($c = 1.0$, CDCl_3), in 67% yield.⁶

In order to demonstrate the synthetic utility of this transformation, the elaboration of cycloadduct **3** to shikimic acid (**4**) was undertaken (Scheme II).⁷ Previous syntheses of racemic shikimic acid and related derivatives have originated from Diels-Alder adducts of furan with acrylonitrile⁸ or methyl acrylate.⁹ Attempted ring opening of imide **3** with LiHMDS proved unsuccessful under a variety of conditions; therefore, the imide was converted to methyl ester **5** via the derived intermediate ethylthio ester [a) LiSEt;¹⁰ b) Cs_2CO_3 , MeOH] in 93% overall yield. In a modification of the procedure of Campbell,^{9c} the product of the LiHMDS

induced ring opening of **5** was trapped *in situ* with TBSOTf to give **6** in 90% yield. Dihydroxylation (OsO_4 , NMO, THF, H_2O) of diene **6** proceeded with good diastereoselectivity (10:1) to deliver diol **7** in 76% isolated yield. Desilylation proceeded in 97% yield to provide *ent*-methyl shikimate. Although the saponification of methyl shikimate is reported to be capricious due to varying levels of aromatized products,⁷ it was found that potassium trimethylsilanolate cleanly afforded *ent*-shikimic acid (**4**),¹¹ which was purified by ion-exchange chromatography (IR-120, H_2O)¹² in 90% yield. This material exhibited spectroscopic data comparable to that of a natural sample, with the exception of the rotation, which was opposite in sign $[\alpha]_{\text{D}}^{25}$ (Aldrich) -150° ($c = 0.80$, MeOH); $[\alpha]_{\text{D}}^{25}$ (synthetic) $+142^\circ$ ($c = 0.80$, MeOH)].

Scheme II^a



^a(a) LiSEt, THF, -20°C ; (b) Cs_2CO_3 , MeOH, 0°C ; (c) LiHMDS, THF, -78 to 0°C , then TBSOTf, 2,6-lutidine, -78°C ; (d) OsO_4 , NMO, THF, H_2O , 0°C ; (e) TBAF, THF; (f) TMSOK, THF, then IR-120, H_2O .

In summary, the unnatural enantiomer of shikimic acid has been synthesized in 7 steps and 37% overall yield from imide **2**. The key transformation was the bis(oxazoline)Cu(II)-catalyzed Diels-Alder reaction of acrylimide **2** with furan, which proceeded with excellent enantioselectivity. This route provides a rapid entry into the asymmetric synthesis of this class of compounds.

Acknowledgment. Support has been provided by the National Institutes of Health and the National Science Foundation. A National Science Foundation Predoctoral Fellowship (D.M.B.) is gratefully acknowledged. NMR spectrometers used in this investigation were purchased with funding from the NIH and NSF.

References and Footnotes

- (a) Evans, D. A.; Miller, S. J.; Lectka, T. *J. Am. Chem. Soc.* **1993**, *115*, 6460-6461. (b) Evans, D. A.; Lectka, T.; Miller, S. J. *Tetrahedron Lett.* **1993**, *34*, 7027-7030. (c) Evans, D. A.; Murry, J. A.; von Matt, P.; Norcross, R. D.; Miller, S. J. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 798-800. (d) Evans, D. A.; Kozlowski, M. C.; Tedrow, J. S. *Tetrahedron Lett.* in press.
- (a) Evans, D. A.; Murry, J. A.; Kozlowski, M. C. *J. Am. Chem. Soc.* **1996**, *118*, 5814-5815. (b) Ref. 1d.
- For reviews, see: (a) Hudlicky, T.; Entwistle, D. A.; Pitzer, K. K.; Thorpe, A. J. *Chem. Rev.* **1996**, *96*, 1195-1220. (b) Vogel, P.; Fattori, D.; Gasparini, F.; Le Drian, C. *Synlett* **1990**, 173-185.
- Corey, E. J.; Loh, T.-P. *Tetrahedron Lett.* **1993**, *34*, 3979-3982.
- Yamamoto, I.; Narasaka, K. *Chemistry Lett.* **1995**, 1129-1130.
- Experimental details are as follows: To a mixture of bis(oxazoline)CuCl₂•CH₂Cl₂ (511 mg, 1 mmol) and AgSbF₆ (687 mg, 2 mmol) was added CH₂Cl₂ (15 mL); the mixture was stirred for 3.5 h in the dark. A solution of imide **3** (2.82 g, 20 mmol) in CH₂Cl₂ (25 mL) was cooled to -78°C , and furan (14.5 ml, 200 mmol) was added. The deep-green catalyst solution was filtered through celite into the reaction over 5 min; an additional 5 ml of CH₂Cl₂ was used to wash the celite. After 42 h, the reaction was quenched by the addition of 5 mL conc. NH₄OH solution. On warming, the mixture was diluted with H₂O and extracted three times with CH₂Cl₂. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*. The resulting crystalline solid was dissolved at room temperature in EtOAc, and hexanes were slowly added. After standing overnight at -20°C , the crystals were collected. Yield = 2.76 g (67%). Diastereomer and enantiomer ratios were determined by NMR and chiral HPLC (Chiracel OD-H, 70:30 hexanes:EtOAc, 0.65 mL/min; t^{R} (major) = 24.8 min, t^{R} (minor) = 30.5 min).
- For a recent review of shikimic acid syntheses, see Campbell, M. M.; Sainsbury, M.; Searle, P. A. *Synthesis* **1993**, 179-193.
- Rajapaksa, D.; Keay, B. A.; Rodrigo, R. *Can. J. Chem.* **1984**, *6*, 826-827.
- (a) Campbell, M. M.; Kaye, A. D.; Sainsbury, M. *Tetrahedron Lett.* **1983**, *24*, 4745-4746. (b) Campbell, M. M.; Kaye, A. D.; Sainsbury, M.; Yavarzadeh, R. *Tetrahedron Lett.* **1984**, *25*, 1629-1630. (c) Campbell, M. M.; Kaye, A. D.; Sainsbury, M.; Yavarzadeh, R. *Tetrahedron* **1984**, *40*, 2461-2470. (d) Campbell, M. M.; Sainsbury, M.; Yavarzadeh, R. *Tetrahedron* **1984**, *40*, 5063-5070.
- Damon, R. E.; Coppola, G. M. *Tetrahedron Lett.* **1988**, *29*, 6141-6144.
- Laganis, E. D.; Chenard, B. L. *Tetrahedron Lett.* **1984**, *25*, 5831-5834.
- Pawlak, J. L.; Berchtold, G. A. *J. Org. Chem.* **1987**, *52*, 1765-1771.

(Received in USA 24 October 1996; accepted 11 November 1996)