


A New Strategy for Extending *N*-Acyl Imides as Chiral Auxiliaries for Aldol and Diels–Alder Reactions: Application to an Enantioselective Synthesis of α -Himachalene**

David A. Evans,* David H. Brown Ripin,
Jeffrey S. Johnson, and Eileen A. Shaughnessy

The development of enantioselective transformations involving the ketone functional group continues to be a problem of general interest in asymmetric synthesis due to the ubiquity of this functionality. While chiral Lewis acids have been employed to promote enantioselective aldol and Diels–Alder reactions of carboxylic acid derivatives and aldehydes, stereocontrol is generally low in reactions of ketones due to the lack of differentiation in Lewis acid coordination to the two nonbonding electron pairs of the carbonyl group.^[1] One strategy for creating a stereochemical bias in ketone-derived enolate–electrophile reactions [Eq. (1)] or in the transformations of α,β -unsaturated ketones such as the Diels–Alder reaction [Eq. (2)] has been developed. In



these processes, a stereogenic center, created by the incorporation of the removable substituent X, is positioned proximal to the trigonal carbon centers undergoing reaction, thus establishing a diastereofacial bias for the transformation. The implementation of this strategy for enolates and enolsilanes of α -trialkylsilyl ketones [Eq. (1), X = SiR₃] has been reported by Enders and Lohray,^[2,3] while Heathcock and Masamune have demonstrated that the aldol and Diels–Alder reactions of α -alkoxyketones [Eqs. (1) and (2), X = OR] are highly diastereoselective.^[4]

Here we define the removable auxiliary X = COX_c (X_c = 4-benzyl-2-oxazolidinon-3-yl), which can be used in the previously described enolate bond constructions and combined with a mild hydrolysis/decarboxylation procedure to extend the utility of β -ketoimides^[5] to the production of enantiopure ketones [Eq. (1)]. We also demonstrate that this strategy is applicable to the implementation of enantioselective Diels–Alder reactions [Eq. (2), X = COX_c].


We have previously demonstrated that the β -ketoimide **1**^[6] undergoes metal ion dependent stereoselective *syn*-^[7] and *anti*-aldol^[6] reactions. In addition, the titanium enolate^[8] derived

[*] Prof. D. A. Evans, D. H. B. Ripin, J. S. Johnson, E. A. Shaughnessy
Department of Chemistry and Chemical Biology, Harvard University
Cambridge, MA 02138 (USA)
Fax: Int. code +(617)495-1460
e-mail: evans@chemistry.harvard.edu

[**] Financial support was provided by the U. S. National Science Foundation and the National Institutes of Health.

from **1** has also been shown to undergo diastereoselective orthoester acylation.^[19]

The focus of the current investigation has been the development of a mild decarboxylation process that might be applied to the enolate-derived reaction products that are prone to epimerization and/or elimination.^[10] A successful solution to this transformation is transesterification with ethanethiolate^[11] followed by silver(I)-promoted hydrolysis/decarboxylation^[12] of the resulting thioester (Scheme 1).



Scheme 1. Thiolate displacement, hydrolysis, and decarboxylation of β -oxoimide **2a**[6]: a) EtSH, KH, THF, 25 °C; b) AgNO₃, 2,6-lutidine, THF/H₂O (5/1), 25 °C. TES = triethylsilyl.


Treatment of **2a** with potassium ethanethiolate afforded a mixture of C_α-diastereomeric β -ketothioesters **3**.^[13] Subjecting thioesters **3** to silver(I)-promoted hydrolysis and subsequent decarboxylation at ambient temperature provided ethyl ketone **4a** over the course of 48 hours. In most instances the reaction time may be reduced without compromising yield by mild heating during the hydrolysis step (70 °C, 3 h). The two steps may be merged into a single-flask operation (see Experimental Section). The decarboxylation of β -ketoimide-derived aldol and orthoester acylation adducts **2a–2f** was investigated using this procedure (Table 1). The mild reaction conditions accommodate the presence of ketal, silyl ether, alkene, and acetate functionalities.

Table 1. One-flask decarboxylation of representative β -ketoimide adducts[a].

Imide	Product	Yield [%][b]	t [h][c]
		98	24
		79 74	24 3[d]
		88	48
		93 92	48 3[d]
		83	48


[a] X_q = (S)-4-benzyl-2-oxazolidinon-3-yl, X_r = (R)-4-benzyl-2-oxazolidinon-3-yl, TBS = *tert*-butyldimethylsilyl. [b] Yield after purification. [c] Reaction time for the hydrolysis step. [d] Hydrolysis run at reflux.

These ketone decarboxylation products are useful building blocks for the rapid synthesis of extended polypropionate chains.^[14] In the example shown in Scheme 2, polypropionate **5**, containing six stereocenters, was synthesized in four steps (63 % yield) from β -ketoimide **1**; all stereocenters were formed with a combined diastereoselectivity exceeding 95%.^[14e]



Scheme 2. Strategy for the rapid assembly of polypropionate chains. a) 1. TiCl₄, Et₃N, 0 °C, 2. *i*PrCHO, -78 °C; b) TBSOTf, 2,6-lutidine, -78 °C; c) 1. EtSH, KH, 2. AgNO₃, 2,6-lutidine, H₂O; d) chex₂BrCl, Et₃N, -78 °C. OTf = trifluoromethanesulfonate, PMP = 4-methoxybenzyl.

This auxiliary-based methodology can be applied to achieve control of the absolute stereochemistry in the Diels–Alder process [Eq. (2)]. In this context, we investigated an intramolecular Diels–Alder reaction that comprises the pivotal bond construction in an enantioselective synthesis of the sesquiterpene α -himachalene (**11**).^[15] Dienimide **6**, prepared in five steps from 3,3-dimethyl-4-propenal, underwent a highly diastereoselective boron-mediated aldol reaction with acrolein to afford the allylic alcohol **7** as the sole isomer (Scheme 3).^[16] Compound **7** was subjected to Parikh–Doering oxidation^[17] and the resulting trienone **8** was treated with ZnBr₂ (0 °C, CH₂Cl₂, 1.5 h) to effect cycloaddition (78 % yield, two steps) to the desired crystalline *endo* adduct **9** (97:3 diastereoselectivity), which was isolated as



Scheme 3. Enantioselective synthesis of α -himachalene (**11**). a) 1. Bu₂BOTf, *i*Pr₂NEt, isoprene, 0 °C, 2. acrolein, -78 °C; b) SO₃, pyridine, DMSO/CH₂Cl₂, *i*Pr₂NEt, -10 °C; c) ZnBr₂, 0 °C; d) LiSEt; e) AgNO₃, 2,6-lutidine, THF/H₂O, 70 °C; f) [Cp₂Ti(μ -Cl)(μ -CH₂)AlMe₂], -40 °C.

a single diastereomer by chromatography. The sense of asymmetric induction in this cycloaddition is consistent with a Lewis acid catalyzed Diels–Alder reaction on the *s-cis* ZnBr₂ complex **8a**.

In analogy to previous experiments, Diels–Alder adduct **9** was transformed to the corresponding β -ketothioester in 90% yield and smoothly decarboxylated (70 °C, 24 h) to afford ketone **10** in 86% yield without epimerization at the ring fusion. The synthesis was completed by treatment of **10** with Tebbe reagent^[18] to afford synthetic α -himachalene (**11**) (92% yield), whose spectroscopic and analytical data are identical in all respects to literature values (¹H NMR, IR, TLC, [α]_D).^[15]

The removable auxiliary X = CO₂ described here should prove useful in enantioselective ketone-based bond constructions. Although the methodology has been highlighted with aldol and Diels–Alder reactions, absolute stereochemical control of other transformations such as the Michael reaction should also be possible. Studies extending the scope of these concepts are currently underway.

Experimental Section

General one-flask procedure for the decarboxylation of β -ketoimides: A stirred suspension of KH (1.2 mmol) in THF (10 mL) under argon was charged with EtSH (1.3 mmol) and gas evolved. The white suspension was stirred for 1 h before a solution of β -oxoimide (1 mmol) in THF (10 mL) was added by cannula. The reaction mixture was stirred until the starting material had been consumed. To the reaction mixture was added water (4 mL) and 2,6-lutidine (10 mmol), followed by AgNO₃ (2.5 mmol). The reaction mixture was protected from light and stirred for two days. The pale yellow suspension was filtered through Celite with ether, and the organic solution was extracted between ether (50 mL) and aqueous CuSO₄ solution (50 mL). The ether layer was separated, dried over MgSO₄, and concentrated in vacuo. Chromatography on silica gel using an appropriate solvent mixture provided pure ketone.

Received: April 7, 1997 [Z 10313 IE]

German version: *Angew. Chem.* 1997, 109, 2208–2210

Keywords: aldol reactions • chiral auxiliaries • decarboxylations • Diels–Alder reactions • himachalene

- [1] S. Shambayati, S. L. Schreiber in *Comprehensive Organic Synthesis*, Vol. 1 (Eds.: B. M. Trost, I. Fleming), Pergamon, New York, 1991, pp. 283–324.
- [2] a) D. Enders, B. B. Lohray, *Angew. Chem.* 1988, 100, 594–596; *Angew. Chem. Int. Ed. Engl.* 1988, 27, 581–582; b) B. B. Lohray, D. Enders, *Helv. Chim. Acta* 1989, 72, 980–984; c) B. B. Lohray, R. Zimbiniski, *Tetrahedron Lett.* 1990, 31, 7273–7276.
- [3] Another important approach to the construction of ketone-derived chiral auxiliaries has been through the use of chiral metalloenamines: a) S. F. Martin in *Comprehensive Organic Synthesis*, Vol. 2 (Eds.: B. M. Trost, I. Fleming), Pergamon, New York, 1991, pp. 475–502.
- [4] a) C. H. Heathcock, M. C. Pirrung, C. T. Buse, J. P. Hagen, S. D. Young, J. E. Sohn, *J. Am. Chem. Soc.* 1979, 101, 7077–7079; b) S. Masamune, W. Choy, F. A. Kerdesky, B. Imperiali, *J. Am. Chem. Soc.* 1981, 103, 1566–1568; c) S. Masamune, L. A. Reed III, J. T. Davis, W. Choy, *J. Org. Chem.* 1983, 48, 1154–1156.
- [5] D. A. Evans, M. D. Ennis, T. Le, N. Mandel, G. Mandel, *J. Am. Chem. Soc.* 1984, 106, 1154–1156.
- [6] The experimental details for the synthesis of **1** have been reported: D. A. Evans, H. P. Ng, J. S. Clark, D. L. Rieger, *Tetrahedron* 1992, 48, 2127–2142.
- [7] D. A. Evans, J. S. Clark, R. Metternich, V. J. Novack, G. S. Sheppard, *J. Am. Chem. Soc.* 1990, 112, 866–868.
- [8] D. A. Evans, F. Urpi, T. C. Somers, J. S. Clark, M. T. Bilodeau, *J. Am. Chem. Soc.* 1990, 112, 8215–8216.
- [9] a) D. A. Evans, H. P. Ng, D. L. Rieger, *J. Am. Chem. Soc.* 1993, 115, 11446–11459 (see the transformation 7 → 18 in Scheme VIII); b) D. A. Evans, A. M. Ratz, B. E. Huff, G. S. Sheppard, *J. Am. Chem. Soc.* 1995, 117, 3448–3467 (see the transformation 4 → 17 in Scheme VII).
- [10] Initial studies utilizing the Krapcho decarboxylation conditions (DMSO, H₂O, 140 °C) on β -oxoimide aldol adducts afforded only moderate yields of the desired ketone products accompanied by significant amounts of β -elimination: a) A. P. Krapcho, *Synthesis* 1982, 805–822, b) 893–914.
- [11] R. E. Damon, G. M. Coppola, *Tetrahedron Lett.* 1990, 31, 2849–2852.
- [12] a) E. J. Corey, M. C. Bock, *Tetrahedron Lett.* 1975, 3269–3270; b) R. Schwyzer, C. Hurlimann, *Helv. Chim. Acta* 1954, 18, 155–166.
- [13] Thiophenolate and 2-methyl-2-propanethiolate both failed to react, while benzylthiolate provides results comparable to those of ethanethiolate.

- [14] a) J. L. Duffy, T. P. Yoon, D. A. Evans, *Tetrahedron Lett.* 1995, 36, 9245–9248; b) D. A. Evans, M. G. Yang, M. J. Dart, J. L. Duffy, *ibid.* 1996, 37, 1957–1960; c) D. A. Evans, M. G. Yang, M. J. Dart, J. L. Duffy, A. S. Kim, *J. Am. Chem. Soc.* 1995, 117, 9598–9599; d) D. A. Evans, D. L. Rieger, M. T. Bilodeau, F. Urpi, *J. Am. Chem. Soc.* 1991, 113, 1047–1049; e) D. A. Evans, M. J. Dart, J. L. Duffy, D. L. Rieger, *ibid.* 1995, 117, 9073–9074.
- [15] a) T. C. Joseph, S. Dev, *Tetrahedron Lett.* 1961, 216–222; b) *Tetrahedron* 1968, 24, 3841–3852, 3853–3859; c) W. Oppolzer, R. L. Snowden, *Helv. Chim. Acta* 1981, 64, 2592–2597; d) E. Wenkert, K. Naemura, *Synth. Comm.* 1973, 3, 45–48.
- [16] D. A. Evans, J. Bartroli, T. L. Shih, *J. Am. Chem. Soc.* 1981, 103, 2127–2129.
- [17] J. R. Parikh, W. von E. Doering, *J. Am. Chem. Soc.* 1967, 89, 5505–5507.
- [18] S. H. Pine, R. Zahler, D. A. Evans, R. H. Grubbs, *J. Am. Chem. Soc.* 1980, 102, 3270–3272.